

Body Mass Index-for-Age Percentiles: Egyptian Boys, 2 to 21 Years

Created from a sample of 13507 Egyptian boys aged 2 to 21 years.

Source: Wright State University School of Medicine Department of Community Health Lifespan Health Research Center in collaboration with the Cairo University Diabetic Endocrine and Pediatric Unit (2002).

Body Mass Index-for-Age Percentiles: Egyptian Girls, 2 to 21 Years



Created from a sample of 13762 Egyptian girls aged 2 to 21 years.

Source: Wright State University School of Medicine Department of Community Health Lifespan Health Research Center in collaboration with the Cairo University Diabetic Endocrine and Pediatric Unit (2002).

Head Circumference-for-Age Percentiles:
Egyptian Boys, Birth to 36 Months



Created from a sample of 3316 Egyptian boys aged birth to 36 months.

Source: Wright State University School of Medicine Department of Community Health Lifespan Health Research Center in collaboration with the Cairo University Diabetic Endocrine and Pediatric Unit (2002).



Head Circumference-for-Age Percentiles: Egyptian Girls, Birth to 36 Months

Created from a sample of 2735 Egyptian girls aged birth to 36 months.

Source: Wright State University School of Medicine Department of Community Health Lifespan Health Research Center in collaboration with the Cairo University Diabetic Endocrine and Pediatric Unit (2002).



Weight-for-Age Percentiles:
Egyptian Boys, Birth to 36 Months

Created from a sample of 3645 Egyptian boys aged birth to 36 months.

Source: Wright State University School of Medicine Department of Community Health Lifespan Health Research Center in collaboration with the Cairo University Diabetic Endocrine and Pediatric Unit (2002).

Weight-for-Age Percentiles: Egyptian Girls, Birth to 36 Months



Created from a sample of 3016 Egyptian girls aged birth to 36 months.

Source: Wright State University School of Medicine Department of Community Health Lifespan Health Research Center in collaboration with the Cairo University Diabetic Endocrine and Pediatric Unit (2002).

Recumbent Length-for-Age Percentiles: Egyptian Boys, Birth to 36 Months



Created from a sample of 3302 Egyptian boys aged birth to 36 months.

Source: Wright State University School of Medicine Department of Community Health Lifespan Health Research Center in collaboration with the Cairo University Diabetic Endocrine and Pediatric Unit (2002).

Recumbent Length-for-Age Percentiles: Egyptian Girls, Birth to 36 Months



Created from a sample of 2770 Egyptian girls aged birth to 36 months.

Source: Wight State University School of Medicine Department of Community Health Lifespan Health Research Center in collaboration with the Cairo University Diabetic Endocrine and Pediatric Unit (2002).



Stature-for-Age Percentiles: Egyptian Boys, 2 to 21 Years

Created from a sample of 13533 Egyptian boys aged 2 to 21 years.

Source: Wright State University School of Medicine Department of Community Health Lifespan Health Research Center in collaboration with the Cairo University Diabetic Endocrine and Pediatric Unit (2002).

Stature-for-Age Percentiles: Egyptian Girls, 2 to 21 Years



Created from a sample of 13809 Egyptian girls aged 2 to 21 years.

Source: Wright State University School of Medicine Department of Community Health Lifespan Health Research Center in collaboration with the Cairo University Diabetic Endocrine and Pediatric Unit (2002).

Weight-for-Age Percentiles: Egyptian Boys, 2 to 21 Years



Created from a sample of 13703 Egyptian boys aged 2 to 21 years.

Source: Wright State University School of Medicine Department of Community Health Lifespan Health Research Center in collaboration with the Cairo University Diabetic Endocrine and Pediatric Unit (2002).

Weight-for-Age Percentiles:
Egyptian Girls, 2 to 21 Years



Created from a sample of 13933 Egyptian girls aged 2 to 21 years.

Source: Wright State University School of Medicine Department of Community Health Lifespan Health Research Center in collaboration with the Cairo University Diabetic Endocrine and Pediatric Unit (2002).



Weight-for-Recumbent Length Percentiles: Egyptian Boys, Birth to 36 Months

Created from a sample of 3068 Egyptian boys aged birth to 36 months.

Source: Wright State University School of Medicine Department of Community Health Lifespan Health Research Center in collaboration with the Cairo University Diabetic Endocrine and Pediatric Unit (2002).

Weight-for-Recumbent Length Percentiles: Egyptian Girls, Birth to 36 Months



Created from a sample of 2602 Egyptian girls aged birth to 36 months.

Source: Wright State University School of Medicine Department of Community Health Lifespan Health Research Center in collaboration with the Cairo University Diabetic Endocrine and Pediatric Unit (2002).